

On the LSL for random fields

Allan Gut
Uppsala University

Ulrich Stadtmüller
University of Ulm

Abstract

In some earlier work we have considered extensions of Lai's (1974) law of the single logarithm for delayed sums to a multiindex setting with the same as well as different expansion rates in the various dimensions. A further generalization concerns window sizes that are regularly varying with index 1 (on the line). In the present paper we establish multiindex versions of the latter as well as for some mixtures of expansion rates. In order to keep things within reasonable size we confine ourselves to some special cases for the index set $\mathbb{Z}_+^2$.

1 Introduction

Let $X, \{X_k, k \geq 1\}$ be i.i.d. random variables with mean $\mu = 0$ and partial sums $\{S_n, n \geq 1\}$. The Hartman-Wintner *Law of the Iterated Logarithm* (LIL) [10] states that

$$\limsup_{n \rightarrow \infty} \frac{S_n}{\sqrt{2n \log \log n}} = \sigma \quad \text{a.s.}$$

Later Strassen, in [16], proved the necessity in the sense that he showed that if

$$P\left(\limsup_{n \rightarrow \infty} \frac{|S_n|}{\sqrt{n \log \log n}} < \infty\right) > 0,$$

then $EX^2 < \infty$ and $EX = 0$.

REMARK 1.1 By the Kolmogorov zero-one law it follows that if the probability of the limsup being finite is positive, then it is automatically equal to 1.

REMARK 1.2 Here and throughout there also exist obvious analogs for the limit inferior. $\square$

The *Law of the Single Logarithm* (LSL) is due to Lai [11], and deals with *delayed sums* or *windows*, viz.,

$$T_{n,n+k} = \sum_{j=n+1}^{n+k} X_j, \quad n \geq 0, k \geq 1,$$

and states that, if

$$E|X|^{2/\alpha} (\log^+ |X|)^{-1/\alpha} < \infty \quad \text{and} \quad EX = 0, EX^2 = \sigma^2,$$

then, for $0 < \alpha < 1$,

$$\limsup_{n \rightarrow \infty} \frac{T_{n,n+n^\alpha}}{\sqrt{2n^\alpha \log n}} = \sigma \sqrt{1-\alpha} \quad \text{a.s.},$$

and, conversely, that if

$$P\left(\limsup_{n \rightarrow \infty} \frac{|T_{n,n+n^\alpha}|}{\sqrt{n^\alpha \log n}} < \infty\right) > 0,$$

AMS 2000 subject classifications. Primary 60F15, 60G50; Secondary 60G70.

Keywords and phrases. Delayed sums, window, law of the iterated logarithm, law of the single logarithm, sums of i.i.d. random variables, window, slowly varying function, multidimensional indices, random fields.

Abbreviated title. On the LSL for random fields.

Date. December 4, 2009

then

$$E|X|^{2/\alpha}(\log^+|X|)^{-1/\alpha} < \infty \quad \text{and} \quad EX = 0.$$

REMARK 1.3 Here, and throughout, $\log^+ x = \max\{\log x, 1\}$ for $x > 0$. $\square$

These results can be generalized in various ways. However, let us first mention that there also exist *one-sided* versions of the above results.

Martikainen [12], Rosalsky [15] and Pruitt [14] independently proved a one-sided LIL to the effect that if

$$-\infty < \limsup_{n \rightarrow \infty} \frac{S_n}{\sqrt{2n \log \log n}} < \infty$$

with positive probability, then $EX^2 < \infty$ and $EX = 0$. An analogous one-sided version of Lai's result is Theorem 3 of [2], where now the finiteness of the one-sided limsup is equivalent to the same moment condition as in [11], however, based on $X^+ = \max\{X, 0\}$ rather than on $|X|$.

As for multiindex results, Wichura [17] proved the following LIL for random fields: Let $\{X_{\mathbf{k}}, \mathbf{k} \in \mathbf{Z}_+^d\}$ be i.i.d. random variables with partial sums $S_{\mathbf{n}} = \sum_{\mathbf{k} \leq \mathbf{n}} X_{\mathbf{k}}$, $\mathbf{n} \in \mathbf{Z}_+^d$, where the random field or index set $\mathbf{Z}_+^d$, $d \geq 2$, is the positive integer d -dimensional lattice with coordinate-wise partial ordering $\leq$, where $|\mathbf{n}| = \prod_{k=1}^d n_k$, and where $\mathbf{n} \rightarrow \infty$ means that $n_k \rightarrow \infty$ for all $k = 1, 2, \dots, d$. Then

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{S_{\mathbf{n}}}{\sqrt{2|\mathbf{n}| \log \log |\mathbf{n}|}} = \sigma\sqrt{d} \quad \text{a.s.},$$

provided

$$EX^2 \frac{(\log^+|X|)^{d-1}}{\log^+ \log^+|X|} < \infty \quad \text{and} \quad EX = 0, \quad EX^2 = \sigma^2,$$

and, conversely, if

$$P\left(\limsup_{\mathbf{n} \rightarrow \infty} \frac{|S_{\mathbf{n}}|}{\sqrt{2|\mathbf{n}| \log \log |\mathbf{n}|}} < \infty\right) > 0,$$

then

$$EX^2 \frac{(\log^+|X|)^{d-1}}{\log^+ \log^+|X|} < \infty \quad \text{and} \quad EX = 0.$$

The analogous multiindex extension of Lai's result is given in [8]. In order to describe the main result there, let $\{X_{\mathbf{k}}, \mathbf{k} \in \mathbf{Z}_+^d\}$ be i.i.d. random variables and define the delayed sums—which in this setting turn into “real” windows,

$$T_{\mathbf{n}, \mathbf{n} + \mathbf{n}^\alpha} = \sum_{\mathbf{k} \leq \mathbf{n} + \mathbf{n}^\alpha} X_{\mathbf{k}}, \quad \mathbf{n} \in \mathbf{Z}_+^d,$$

and where addition is to be taken coordinate-wise. Then, for $0 < \alpha < 1$,

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{T_{\mathbf{n}, \mathbf{n} + \mathbf{n}^\alpha}}{\sqrt{2|\mathbf{n}|^\alpha \log |\mathbf{n}|}} = \sigma\sqrt{1-\alpha} \quad \text{a.s.},$$

provided

$$E|X|^{2/\alpha}(\log^+|X|)^{d-1-1/\alpha} < \infty \quad \text{and} \quad EX = 0, \quad EX^2 = \sigma^2.$$

We also proved that if

$$P\left(\limsup_{\mathbf{n} \rightarrow \infty} \frac{|T_{\mathbf{n}, \mathbf{n} + \mathbf{n}^\alpha}|}{\sqrt{|\mathbf{n}|^\alpha \log |\mathbf{n}|}} < \infty\right) > 0,$$

then

$$E|X|^{2/\alpha}(\log^+|X|)^{d-1-1/\alpha} < \infty \quad \text{and} \quad EX = 0.$$

Note in particular that the moment condition depends on the d (as in Wichura's LIL above).

In [9] this result was generalized to allow for different α 's for the different directions, and it was shown there that if

$$E|X|^{2/\alpha_1}(\log^+|X|)^{p-1-1/\alpha_1} < \infty \quad \text{and} \quad EX = 0, \quad EX^2 = \sigma^2,$$

then

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{T_{\mathbf{n}, \mathbf{n} + \mathbf{n}^\alpha}}{\sqrt{2|\mathbf{n}^\alpha| \log |\mathbf{n}|}} = \sigma \sqrt{1 - \alpha_1} \quad \text{a.s.},$$

and, conversely, that if

$$P\left(\limsup_{\mathbf{n} \rightarrow \infty} \frac{|T_{\mathbf{n}, \mathbf{n} + \mathbf{n}^\alpha}|}{\sqrt{|\mathbf{n}^\alpha| \log |\mathbf{n}|}} < \infty\right) > 0,$$

then

$$E|X|^{2/\alpha_1}(\log^+ |X|)^{p-1-1/\alpha_1} < \infty \quad \text{and} \quad EX = 0,$$

where now

$$0 < \alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_d < 1, \quad p = \max\{k : \alpha_k = \alpha_1\} \quad \text{and} \quad |\mathbf{n}^\alpha| = \prod_{k=1}^d n_k^{\alpha_k}.$$

REMARK 1.4 When $\alpha_1 = \alpha_2 = \dots = \alpha_d = \alpha$ the theorem reduces (of course) to the previous one. $\square$

The next natural question is to consider the boundary cases $\alpha = 0$ and $\alpha = 1$. As for the former, one case is the trivial one in which the windows reduce to single random variables, viz., $T_{n, n+1} = X_{n+1}$. A nondegenerate variation concerns the delayed sums $T_{n, n+\log n}$, $n \geq 1$, which obey the so-called Erdős-Rényi law ([4], Theorem 2, [3], Theorem 2.4.3).

In [7] we considered the nondegenerate boundary case $\alpha = 1$ at the other end. In this case the window size is larger than any power less than one, and at the same time not quite linear. Technically, the paper focused on windows of the form

$$T_{n, n+a_n} \quad \text{where} \quad a_n = \frac{n}{L(n)} \quad \text{with} \quad (1.1)$$

$$\text{a differentiable function} \quad L(\cdot) \nearrow \infty \in \mathcal{SV} \quad \text{and} \quad \frac{xL'(x)}{L(x)} \searrow \quad \text{as} \quad x \rightarrow \infty, \quad (1.2)$$

where $L \in \mathcal{SV}$ means that L is slowly varying at infinity (see e.g. [1] or [6], Section A.7), and where, for convenience, we shall permit ourselves to treat quantities such as $a_n = n/L(n)$ and $a_n = n/\log n$ and so on as integers.

Finally, let

$$d_n = \log \frac{n}{a_n} + \log \log n = \log L(n) + \log \log n, \quad n \geq 2,$$

and set

$$f_n = \min\{a_n \cdot d_n, n\},$$

with f an increasing interpolating function, i.e., $f(x) = f_{[x]}$ for $x > 0$, and with f^{-1} the corresponding (suitably defined) inverse function. Then, in short, (the precise equivalence has to be formulated as above)

$$\limsup_{n \rightarrow \infty} \frac{T_{n, n+a_n}}{\sqrt{2a_n d_n}} = \sigma \quad \text{a.s.} \quad \iff \quad E(f^{-1}(X^2)) < \infty, \quad EX = 0, \quad EX^2 = \sigma^2.$$

As an introduction and point of departure to what follows, here are the two canonical examples; [7], Corollaries 2.1 and 2.2, which concern the cases $L(n) = \log n$ and $L(n) = \log \log n$, that is,

$$d_n = 2 \log \log n, \quad f_n = 2 \frac{n}{\log n} \log \log n,$$

and

$$d_n = \log \log \log n + \log \log n \sim \log \log n, \quad f_n \sim n,$$

respectively.

Theorem 1.1 Let $X, \{X_k, k \geq 1\}$ be i.i.d. random variables with partial sums $S_n = \sum_{k=1}^n X_k$, $n \geq 1$.

(i) If

$$EX^2 \frac{\log^+ |X|}{\log^+ \log^+ |X|} < \infty \quad \text{and} \quad EX = 0, \quad \text{Var } X = \sigma^2,$$

then

$$\limsup_{n \rightarrow \infty} \frac{T_{n, n+n/\log n}}{\sqrt{4 \frac{n}{\log n} \log \log n}} = \sigma \quad a.s.$$

Conversely, if

$$P\left(\limsup_{n \rightarrow \infty} \frac{|T_{n, n+n/\log n}|}{\sqrt{4 \frac{n}{\log n} \log \log n}} < \infty\right) > 0,$$

then $E X^2 \frac{\log^+ |X|}{\log^+ \log^+ |X|} < \infty$ and $E X = 0$.

(ii) If $E X = 0$ and $E X^2 = \sigma^2 < \infty$, then

$$\limsup_{n \rightarrow \infty} \frac{T_{n, n+n/\log \log n}}{\sqrt{2n}} = \sigma \quad a.s.$$

Conversely, if

$$P\left(\limsup_{n \rightarrow \infty} \frac{|T_{n, n+n/\log \log n}|}{\sqrt{n}} < \infty\right) > 0,$$

then $E X^2 < \infty$ and $E X = 0$. □

The purpose of the present paper is to investigate random field analogs of this result. However, in order to keep things within reasonable bounds we shall confine ourselves to the case $d = 2$ and the windows

$$T_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})} = \sum_{\substack{m \leq i \leq m+a_m^{(1)} \\ n \leq j \leq n+a_n^{(2)}}} X_{i,j}, \quad (1.3)$$




Figure 1: A typical window.

where

1. $a_m^{(1)} = m/\log m$ and $a_n^{(2)} = n/\log n$,
2. $a_m^{(1)} = m/\log \log m$ and $a_n^{(2)} = n/\log \log n$,
3. $a_m^{(1)} = m/\log m$ and $a_n^{(2)} = n/\log \log n$,

leaving further cases to the reader(s).

In Section 2 we present our setup and main result, after which some preliminaries are given in Section 3. Following this, we provide proofs in Section 4. In Section 5 we connect our earlier papers in the area in that we consider windows of the form

$$T_{(m,n), (m+m^\alpha, n+n/\log n)}.$$

Some final remarks, preceding an Appendix in which we have collected some technicalities, are given in Section 6.

2 Setup and main result

In the remainder of the paper we thus suppose that X and $\{X_{i,j}, i, j \geq 1\}$ are i.i.d. random variables with partial sums $S_{m,n} = \sum_{(i,j)=(1,1)}^{(m,n)} X_{i,j}$ and windows $T_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})}$ (see (1.3)) for $m, n \geq 1$.

Theorem 2.1 (i) *If*

$$E X^2 \frac{(\log^+ |X|)^3}{\log^+ \log^+ |X|} < \infty \quad \text{and} \quad E X = 0, \quad E X^2 = \sigma^2,$$

then

$$\limsup_{m,n \rightarrow \infty} \frac{T_{(m,n), (m+m/\log m, n+n/\log n)}}{\sqrt{4mn \frac{\log \log m + \log \log n}{\log m \log n}}} = \sigma \quad a.s.$$

Conversely, if

$$P\left(\limsup_{n \rightarrow \infty} \frac{|T_{(m,n), (m+m/\log m, n+n/\log n)}|}{\sqrt{mn \frac{\log \log m + \log \log n}{\log m \log n}}} < \infty\right) > 0,$$

then $E X^2 \frac{(\log^+ |X|)^3}{\log^+ \log^+ |X|} < \infty$ *and* $E X = 0$.

(ii) *If*

$$E X^2 \log^+ |X| \log^+ \log^+ |X| < \infty \quad \text{and} \quad E X = 0, \quad E X^2 = \sigma^2,$$

then

$$\limsup_{m,n \rightarrow \infty} \frac{T_{(m,n), (m+m/\log \log m, n+n/\log \log n)}}{\sqrt{2mn \frac{\log \log m + \log \log n}{\log \log m \log \log n}}} = \sigma \quad a.s.$$

Conversely, if

$$P\left(\limsup_{n \rightarrow \infty} \frac{|T_{(m,n), (m+m/\log \log m, n+n/\log \log n)}|}{\sqrt{mn \frac{\log \log m + \log \log n}{\log \log m \log \log n}}} < \infty\right) > 0,$$

then $E X^2 \log^+ |X| \log^+ \log^+ |X| < \infty$ *and* $E X = 0$.

(iii) *If*

$$E X^2 (\log^+ |X|)^2 < \infty \quad \text{and} \quad E X = 0, \quad E X^2 = \sigma^2,$$

then

$$\limsup_{m,n \rightarrow \infty} \frac{T_{(m,n), (m+m/\log m, n+n/\log \log n)}}{\sqrt{4mn \frac{\log \log m + \log \log n}{\log m \log \log n}}} = \sigma \quad a.s.$$

Conversely, if

$$P\left(\limsup_{n \rightarrow \infty} \frac{|T_{(m,n), (m+m/\log m, n+n/\log \log n)}|}{\sqrt{4mn \frac{\log \log m + \log \log n}{\log m \log \log n}}} < \infty\right) > 0,$$

then $E X^2 (\log^+ |X|)^2 < \infty$ *and* $E X = 0$.

3 Preliminaries

Before we jump into the proof of Theorem 2.1 we present the Kolmogorov exponential bounds (cf. e.g. [6], Section 8.2) adapted to the present situation, a lemma that relates certain tail sums to moments together with some special cases that will be of use later.

But first some notation. Let, for $i = 1, 2$ and $n \geq 2$,

$$d_n^{(i)} = \log \frac{n}{a_n^{(i)}} + \log \log n = \log L_i(n) + \log \log n \quad \text{and} \quad r_n = \max\{d_n^{(1)}, d_n^{(2)}\}.$$

The two-dimensional analogs of a_n , d_n , and f_n are as follows:

$$a_{m,n} = a_m^{(1)} \cdot a_n^{(2)} = \frac{m}{L_1(m)} \cdot \frac{n}{L_2(n)}, \quad d_{m,n} = r_m + r_n, \quad f_{m,n} = a_{m,n} \cdot d_{m,n},$$

where in the following the slowly varying functions L are logarithms or iterated logarithms.

In order to introduce the traditional double-truncation we let $\delta > 0$ and $\varepsilon > 0$, and set

$$b_{m,n} = \frac{\sigma\delta}{\varepsilon} \sqrt{\frac{a_{m,n}}{d_{m,n}}}, \quad (3.1)$$

and note that

$$\frac{a_{m,n}}{b_{m,n}} = \frac{\varepsilon}{\sigma\delta} \sqrt{f_{m,n}}. \quad (3.2)$$

Next we set

$$\begin{aligned} X'_{m,n} &= X_{m,n} I\{|X_{m,n}| \leq b_{m,n}\}, & X''_{m,n} &= X_{m,n} I\{b_{m,n} < |X_{m,n}| < \delta\sqrt{f_{m,n}}\}, \\ X'''_{m,n} &= X_{m,n} I\{|X_{m,n}| \geq \delta\sqrt{f_{m,n}}\}. \end{aligned}$$

Throughout the following all objects with primes or multiple primes refer to the respective truncated summands.

3.1 Exponential bounds

The standard procedure for estimating $E X'_{i,j}$ yields

$$|E X'_{i,j}| \leq E|X| I\{|X| > b_{i,j}\} \leq \frac{E X^2 I\{|X| > b_{i,j}\}}{b_{i,j}},$$

so that, omitting intermediate steps and recalling (3.2),

$$\begin{aligned} |E T'_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})}| &\leq a_{m,n} \frac{E X^2 I\{|X| > b_{m,n}\}}{b_{m,n}} \\ &= \frac{\varepsilon}{\sigma\delta} \sqrt{f_{m,n}} E X^2 I\{|X| > b_{m,n}\} \\ &= o(\sqrt{f_{m,n}}) \quad \text{as } m, n \rightarrow \infty. \end{aligned}$$

Moreover, a simple calculation yields, for any $\delta > 0$ and $\max\{m, n\} \geq n_0(\delta)$,

$$a_{m,n} \sigma^2 (1 - \delta) \leq \text{Var} (T'_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})}) \leq a_{m,n} \sigma^2.$$

Inserting the above estimates into the Kolmogorov exponential bounds then yields

$$P\left(T'_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})} > \varepsilon \sqrt{2f_{m,n}}\right) \begin{cases} \leq \exp\left\{-\frac{\varepsilon^2(1-\delta)^3}{\sigma^2} \cdot d_{m,n}\right\}, \\ \geq \exp\left\{-\frac{\varepsilon^2(1+\delta)^2(1+\gamma)}{\sigma^2(1-\delta)} \cdot d_{m,n}\right\}, \end{cases} \quad (3.3)$$

for any $\gamma > 0$ and $\max\{m, n\}$ large.

3.2 A useful lemma

The present subsection contains a technical lemma and some consequences that will be of use later.

Lemma 3.1 *Consider a positive function G on $[1, \infty) \times [1, \infty]$, such that, for some constants $0 < c_1 \leq c_2$ and for all $m, n \in \mathbf{N}$,*

$$c_1 \leq \frac{G(m, n)}{G(u, v)} \leq c_2 \quad \text{for } (u, v) \in [m, m+1] \times [n, n+1]. \quad (3.4)$$

Furthermore, let

$$M(x) = |\{(u, v) \in [1, \infty) \times [1, \infty) : G(u, v) \leq x\}|, \quad (3.5)$$

and assume that, for any $c > 1$, the quotient $\frac{M(cx)}{M(x)}$ is bounded as $x \rightarrow \infty$. Finally, let X be a nonnegative random variable. Then

$$\sum_{m,n=1}^{\infty} P(X > G(m, n)) < \infty \iff \iint P(X > G(u, v)) du dv < \infty \iff E M(X) < \infty.$$

PROOF. The equivalences follow from

$$\sum_{m,n=1}^{\infty} P(X > G(m,n)) = \sum_{m,n=1}^{\infty} \int_m^{m+1} \int_n^{n+1} P\left(X > \frac{G(m,n)}{G(u,v)} G(u,v)\right) dudv,$$

(3.4), and the fact that

$$\begin{aligned} \iint P(X > G(u,v)) dudv &= \iint \left(\int_{G(u,v)}^{\infty} dF_X(x) \right) du dv \\ &= \int_0^{\infty} \left(\iint_{\{(u,v): G(u,v) \leq x\}} dudv \right) dF_X(x) \\ &= \int_0^{\infty} M(x) dF_X(x) = E M(X). \end{aligned} \quad \square$$

Next, some useful examples that we collect separately. The first part is immediate from the lemma and the Borel–Cantelli lemmas. For the verification of the special cases we refer to the appendix.

Corollary 3.1 *Suppose that $\{X, X_{m,n}, m, n \geq 1\}$ are i.i.d. random variables and let M and G be defined as in the lemma. Then*

$$\limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{G(m,n)}} < \infty \quad a.s. \iff \sum_{m,n} P(|X| > \sqrt{G(m,n)}) < \infty \iff E M(X^2) < \infty.$$

In particular,

$$\begin{aligned} \limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{mn \frac{\log \log m + \log \log n}{\log m \log n}}} < \infty \quad a.s. &\iff \sum_{m,n} P\left(|X| > \sqrt{mn \frac{\log \log m + \log \log n}{\log m \log n}}\right) < \infty \\ &\iff E X^2 \frac{(\log^+ |X|)^3}{\log^+ \log^+ |X|} < \infty, \end{aligned} \quad (3.6)$$

$$\begin{aligned} \limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{mn \frac{\log \log m + \log \log n}{\log \log m \log \log n}}} < \infty \quad a.s. &\iff \sum_{m,n} P\left(|X| > \sqrt{mn \frac{\log \log m + \log \log n}{\log \log m \log \log n}}\right) < \infty \\ &\iff E X^2 \log^+ |X| \log^+ \log^+ |X| < \infty, \end{aligned} \quad (3.7)$$

$$\begin{aligned} \limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{mn \frac{\log \log m + \log \log n}{\log m \log \log n}}} < \infty \quad a.s. &\iff \sum_{m,n} P\left(|X| > \sqrt{mn \frac{\log \log m + \log \log n}{\log m \log \log n}}\right) < \infty \\ &\iff E X^2 (\log^+ |X|)^2 < \infty. \end{aligned} \quad (3.8)$$

4 Proof of Theorem 2.1

Since the proof follows the pattern of our predecessors we confine ourselves to providing the proof of part (i) of the theorem in somewhat more detail and the other cases more sketchily.

4.1 Proof of Theorem 2.1(i)

Exponential bounds

In this case we have

$$a_{m,n} = \frac{mn}{\log m \log n}, \quad d_{m,n} = 2(\log \log m + \log \log n), \quad f_{m,n} = \frac{2mn}{\log m \log n} \cdot (\log \log m + \log \log n),$$

so that the Kolmogorov exponential bounds (3.3) yield

$$P\left(T'_{(m,n), (m+m/\log m, n+n/\log n)} > \varepsilon \sqrt{2 \frac{mn(\log \log m + \log \log n)}{\log m \log n}}\right) \begin{cases} \leq \exp\left\{-\frac{\varepsilon^2(1-\delta)^3}{\sigma^2} \cdot 2(\log \log m + \log \log n)\right\}, \\ \geq \exp\left\{-\frac{\varepsilon^2(1+\delta)^2(1+\gamma)}{\sigma^2(1-\delta)} \cdot 2(\log \log m + \log \log n)\right\}, \end{cases} \quad (4.1)$$

for any $\gamma > 0$ and $\max\{m, n\}$ large.

Upper bound

In order to obtain an upper bound we consider, for some constant $c > 0$ to be chosen later, the subset

$$(m_i, n_j) = (e^{\sqrt{ci}}, e^{\sqrt{cj}}) \quad \text{for } i, j \geq \max\{1/c, 1\}. \quad (4.2)$$

The upper exponential bound then reduces to

$$P\left(T'_{(m_i, n_j), (m_i+m_i/\log m_i, n_j+n_j/\log n_j)} > \varepsilon \sqrt{2 \cdot \frac{mn(\log \log m_i + \log \log n_j)}{\log m_i \log n_j}}\right) \leq C \exp\left\{-\frac{\varepsilon^2(1-\delta)^3}{\sigma^2} \cdot (\log i + \log j)\right\} = C(ij)^{-\frac{\varepsilon^2(1-\delta)^3}{\sigma^2}}, \quad (4.3)$$

from which it follows that

$$\sum_{i,j} P\left(T'_{(m_i, n_j), (m_i+m_i/\log m_i, n_j+n_j/\log n_j)} > \varepsilon \sqrt{2 \frac{m_i n_j (\log \log m_i + \log \log n_j)}{\log m_i \log n_j}}\right) < \infty \quad (4.4)$$

whenever $\varepsilon > \sigma(1-\delta)^{-3/2}$.

As for the contribution of T'' we observe as in [8] that in order for $|T''_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})}|$ to surpass the level $\eta\sqrt{f_{m,n}}$ it is necessary that at least $N \geq \eta/\delta$ of the X'' :s are nonzero, which, by stretching the truncation bounds to the extremes, implies that

$$\begin{aligned} P(|T''_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})}| > \eta\sqrt{f_{m,n}}) &\leq \binom{a_m^{(1)} a_n^{(2)}}{N} \left(P(b_{m,n} < |X| \leq \delta\sqrt{f_{m,n}})\right)^N \\ &\leq C \left(a_m^{(1)} a_n^{(2)}\right)^N \left(\frac{E H(|X|)}{H(b_{m,n})}\right)^N, \end{aligned} \quad (4.5)$$

where $E H(|X|) < \infty$ is the appropriate moment condition.

In the present case this amounts, after simplifying, to

$$\begin{aligned} P''(m, n) &= P(|T''_{(m,n), (m+m/\log m, n+n/\log n)}| > \eta\sqrt{f_{m,n}}) \\ &\leq C \left(\frac{(\log \log m + \log \log n)(\log \log(mn))}{(\log(mn))^3}\right)^N. \end{aligned} \quad (4.6)$$

This means that for our subset (4.2) we have

$$P''(m_i, n_j) \leq C \left(\frac{(\log(ci) + \log(cj))(\log(\sqrt{ci} + \sqrt{cj}))}{(\sqrt{ci} + \sqrt{cj})^3}\right)^N \leq C \left(\frac{(\log i + \log j)(\log(i+j))}{i^{3/2} + j^{3/2}}\right)^N. \quad (4.7)$$

Since the sum of these probabilities converges whenever $N \geq 2$, we conclude, via the first Borel–Cantelli lemma, considering in addition that $N\delta \geq \eta$, that

$$\limsup_{i,j \rightarrow \infty} \frac{|T''_{(m_i, n_j), (m_i+m_i/\log m_i, n_j+n_j/\log n_j)}|}{\sqrt{f_{m_i, n_j}}} \leq \delta \quad \text{a.s.} \quad (4.8)$$

The next step is to show that

$$\lim_{m,n \rightarrow \infty} \frac{T'''_{(m,n), (m+m/\log m, n+n/\log n)}}{\sqrt{f_{m,n}}} = 0 \quad \text{a.s.} \quad (4.9)$$

Now, since in order for $|T'''_{(m,n), (m+m/\log m, n+n/\log n)}|$ to surpass the level $\eta\sqrt{f_{m,n}}$ infinitely often it is necessary that infinitely many of the X''' 's are nonzero. However, via an appeal to the first Borel–Cantelli lemma, the latter event has zero probability. Namely, for every $\eta > 0$ we have

$$\sum_{m,n} P(|X_{m,n}| > \eta\sqrt{f_{m,n}}) = \sum_{m,n} P\left(|X| > \eta\sqrt{\frac{2mn}{\log m \log n}} \cdot (\log \log m + \log \log n)\right) < \infty,$$

since, by assumption, $E X^2 \frac{(\log^+ |X|)^3}{\log^+ \log^+ |X|} < \infty$; recall (3.6).

By combining (4.4), (4.8), and (4.9) we are now in the position to conclude that

$$\limsup_{i,j \rightarrow \infty} \frac{T_{(m_i, n_j), (m_i+m_i/\log m_i, n_j+n_j/\log n_j)}}{\sqrt{2f_{m_i, n_j}}} \leq \sigma(1-\delta)^{-3/2} + \delta \quad \text{a.s.}, \quad (4.10)$$

for $(m_i, n_j) = (e^{\sqrt{ci}}, e^{\sqrt{cj}})$ with $i, j \geq 1/c$ and any $c > 0$.

Sufficiency for the entire field

It remains to show that our process behaves accordingly for the entire field. Assume for the moment that the random variables are symmetric, let $\eta > 0$, small, be given, and choose $c = \eta^2$. Recalling that $a_n^{(1)} = a_n^{(2)} = n/\log n$, and that $f_{m,n} = 2mn \frac{\log \log m + \log \log n}{\log m \log n}$, we thus consider

$$\begin{aligned} & P\left(\max_{\substack{m_i \leq m \leq m_{i+1} \\ n_j \leq n \leq n_{j+1}}} \frac{T_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})}}{\sqrt{2 \cdot f_{m,n}}} > (1+12\eta)\sigma\right) \\ & \leq P\left(\max_{\substack{m_i \leq m \leq m_{i+1} \\ n_j \leq n \leq n_{j+1}}} (-T_{(m_i, n_j), (m,n)}) > 2\eta\sigma\sqrt{2f_{m_i, n_j}}\right) \\ & \quad + P\left(\max_{\substack{m_i \leq m \leq m_{i+1} \\ n_j \leq n \leq n_{j+1}}} (-T_{(m_i, n_j), (m+a_m^{(1)}, n)}) > 2\eta\sigma\sqrt{2f_{m_i, n_j}}\right) \\ & \quad + P\left(\max_{\substack{m_i \leq m \leq m_{i+1} \\ n_j \leq n \leq n_{j+1}}} (-T_{(m_i, n_j), (m, n+a_n^{(2)})}) > 2\eta\sigma\sqrt{2f_{m_i, n_j}}\right) \\ & \quad + P\left(\max_{\substack{m_i \leq m \leq m_{i+1} \\ n_j \leq n \leq n_{j+1}}} T_{(m_i, n_j), (m+a_m^{(1)}, n+a_n^{(2)})} > (1+6\eta)\sigma\sqrt{2f_{m_i, n_j}}\right), \end{aligned}$$

where $(m_i, n_j) = (e^{\sqrt{ci}}, e^{\sqrt{cj}})$ as before. Now, since $T \stackrel{d}{=} -T$ the second Lévy inequality for random fields ([13], Theorem 1) is instrumental and we proceed.

$$\begin{aligned} & P\left(\max_{\substack{m_i \leq m \leq m_{i+1} \\ n_j \leq n \leq n_{j+1}}} \frac{T_{(m,n), (m+a_m^{(1)}, n+a_n^{(2)})}}{\sqrt{2f_{m,n}}} > (1+12\eta)\sigma\right) \\ & \leq 2P\left(T_{(m_i, n_j), (m_{i+1}, n_{j+1})} > 2\eta\sigma\sqrt{2f_{m_i, n_j}}\right) \\ & \quad + 2P\left(T_{(m_i, n_j), (m_{i+1}+a_{m_{i+1}}^{(1)}, n_{j+1})} > 2\eta\sigma\sqrt{2f_{m_i, n_j}}\right) \\ & \quad + 2P\left(T_{(m_i, n_j), (m_{i+1}, n_{j+1}+a_{n_{j+1}}^{(2)})} > 2\eta\sigma\sqrt{2f_{m_i, n_j}}\right) \\ & \quad + 2P\left(T_{(m_i, n_j), (m_{i+1}+a_{m_{i+1}}^{(1)}, n_{j+1}+a_{n_{j+1}}^{(2)})} > (1+6\eta)\sigma\sqrt{2f_{m_i, n_j}}\right), \end{aligned}$$

By the definition of our subset we have, for all $\eta > 0$ and $i, j \geq i_0(\eta)$ with some integer $i_0 \geq 1/c$, that (note $c = \eta^2$),

$$\begin{cases} m_{i+1} - m_i \leq e^{\sqrt{ci}} (e^{\sqrt{ci}/(2i)} - 1) \leq \eta^2 a_{m_i}^{(1)}, \\ n_{j+1} - n_j \leq e^{\sqrt{cj}} (e^{\sqrt{cj}/(2j)} - 1) \leq \eta^2 a_{n_j}^{(2)}, \\ a_{m_{i+1}}^{(1)}/a_{m_i}^{(1)} \leq \exp \sqrt{c/i} \leq 1 + 2\eta^2, \\ a_{n_{j+1}}^{(2)}/a_{n_j}^{(2)} \leq \exp \sqrt{c/j} \leq 1 + 2\eta^2, \end{cases} \quad (4.11)$$

from which it follows that the variances satisfy

$$\begin{cases} \text{Var}(T_{(m_i, n_j), (m_{i+1}, n_{j+1})}) \leq \sigma^2 \eta^4 a_{m_i}^{(1)} a_{n_j}^{(2)} = \sigma^2 \eta^4 a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1} + a_{m_{i+1}}^{(1)}, n_{j+1})}) \leq \sigma^2 (1 + 3\eta^2) \eta^2 a_{m_i}^{(1)} a_{n_j}^{(2)} = \sigma^2 (1 + 3\eta^2) \eta^2 a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1}, n_{j+1} + a_{n_{j+1}}^{(2)})}) \leq \sigma^2 (1 + 3\eta^2) \eta^2 a_{m_i}^{(1)} a_{n_j}^{(2)} = \sigma^2 (1 + 3\eta^2) \eta^2 a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1} + a_{m_{i+1}}^{(1)}, n_{j+1} + a_{n_{j+1}}^{(2)})}) \leq \sigma^2 (1 + 3\eta^2)^2 a_{m_i}^{(1)} a_{n_j}^{(2)} = \sigma^2 (1 + 3\eta^2)^2 a_{m_i, n_j}, \end{cases} \quad (4.12)$$

for $i, j \geq i_0$.

From here on the procedure from the previous sections applies for these terms. For example, the second last probability can be bounded with respect to T' by

$$\begin{aligned} 2P\left(T'_{(m_i, n_j), (m_i + \eta^4 a_{m_i}^{(1)}, n_j + (1 + 3\eta^2) a_{n_j}^{(2)})} > 2\eta\sigma\sqrt{2f_{m_i, n_j}}\right) \\ \leq \exp\left\{-\frac{(1 - \delta)^3 4\eta^2 \sigma^2 2 \log(\log m_i \log n_j)}{\sigma^2 \eta^2 (1 + 3\eta^2)}\right\} \leq C(ij)^{-4(1 - \delta)^3}, \end{aligned}$$

provided $\eta^2 < 1/3$, and the last term by

$$\begin{aligned} 2P\left(T'_{(m_i, n_j), (m_i + (1 + 2\eta^2) a_{m_i}^{(1)}, n_j + (1 + 2\eta^2) a_{n_j}^{(2)})} > 2\eta\sigma\sqrt{2f_{m_i, n_j}}\right) \\ \leq \exp\left\{-\frac{(1 - \delta)^3 (1 + 6\eta)^2 \sigma^2 2 \log(\log m_i \log n_j)}{\sigma^2 (1 + 3\eta^2)^2}\right\} \leq C(ij)^{-((1 + 6\eta)/(1 + 3\eta^2))^2 (1 - \delta)^3}. \end{aligned}$$

Both cases yield summable double sequences if δ and η are small enough.

Since constants are not relevant for T'', T''' we finally obtain

$$\sum_{k, j \geq i_0} P\left(\max_{\substack{m_i \leq m \leq m_{i+1} \\ n_j \leq n \leq n_{j+1}}} \frac{T_{(m, n), (m + a_m^{(1)}, n + a_n^{(2)})}}{\sqrt{2f_{m, n}}} > (1 + 12\eta)\sigma\right) < \infty,$$

which implies that the $\limsup \dots \leq (1 + 12\eta)\sigma$, and since η can be chosen arbitrarily small the upper bound is proved for the entire field.

Desymmetrization follows along the usual arguments; note that

$$\frac{E\left|T_{(m, n), (m + a_m^{(1)}, n + a_n^{(2)})}\right|}{\sqrt{f_{m, n}}} \leq \sqrt{\frac{E(T_{(m, n), (m + a_m^{(1)}, n + a_n^{(2)})})^2}{f_{m, n}}} = o(1) \quad \text{as } m, n \rightarrow \infty.$$

Lower bound

Let $c > 2$ and define $m_i = n_i = \exp\{\sqrt{ci}\}$. We first note that

$$\frac{e^{\sqrt{ci}} + e^{\sqrt{ci}/\sqrt{ci}}}{e^{\sqrt{c(i+1)}}} = \frac{1 + 1/\sqrt{ci}}{\exp(\sqrt{ci}(\sqrt{1 + 1/i} - 1))} \sim \frac{1 + 1/\sqrt{ci}}{1 + \sqrt{c/4i}} < 1,$$

eventually and, hence, that the windows of the subset (m_i, n_j) , for i, j large, are disjoint, which means that different blocks (eventually) consist of independent random variables. Using the lower exponential

bound in (4.1) we then obtain

$$\begin{aligned} P\left(T'_{(m_i, n_j)}, (m_i + m_i / \log m_i, n_j + n_j / \log n_j) > (1 - \eta) \sigma \sqrt{2f_{m_i, n_j}}\right) \\ \geq \exp \left\{ \frac{(1 - \eta)^2 (1 + \delta)^2 (1 + \gamma)}{(1 - \delta)} \cdot (2 \log \log m_i + 2 \log \log n_j) \right\} \\ \geq C (ij)^{-\frac{(1 - \eta)^2 (1 + \delta)^2 (1 + \gamma)}{(1 - \delta)}}, \end{aligned}$$

which is a divergent minorant for a choice of $\eta > 0$ such that $1 - \eta < \sqrt{\frac{1 - \delta}{(1 + \delta)^2 (1 + \gamma)}}$. Now, since δ and γ can be chosen arbitrarily small the same is true for η . The desired result finally follows via the Borel–Cantelli Lemma and the fact that T'' and T''' are small.

Strictly speaking, this provides the desired lower bound for our *subset*, after which the overall lower bound follows from that fact that the limsup over *all* windows is at least as large as the limsup over a subset.

Necessity

It follows from the assumption that

$$\limsup_{m, n \rightarrow \infty} \frac{|X_{m, n}|}{\sqrt{f_{m, n}}} = \limsup_{m, n \rightarrow \infty} \frac{|X_{m, n}|}{\sqrt{\frac{2mn(\log \log m + \log \log n)}{\log m \log n}}} < \infty \quad \text{a.s.},$$

from which the necessity of the moment assumption is immediate in view of (3.6). An application of the sufficiency and the strong law of large numbers then tells us that the mean must be equal to zero.

4.2 Proof of Theorem 2.1(ii)

Sufficiency

The proof follows the same procedure with obvious modifications. We have

$$a_{m, n} = \frac{mn}{\log \log m \log \log n}, \quad d_{m, n} = \log \log m + \log \log n, \quad f_{m, n} = \frac{mn(\log \log m + \log \log n)}{\log \log m \log \log n},$$

and the Kolmogorov exponential bounds (3.3) yield

$$\begin{aligned} P(T'_{(m, n)}, (m + m / \log \log m, n + n / \log \log n) > \varepsilon \sqrt{2mn \frac{\log \log m + \log \log n}{\log \log m \log \log n}} \\ \left\{ \begin{aligned} &\leq \exp \left\{ -\frac{\varepsilon^2 (1 - \delta)^3}{\sigma^2} \cdot (\log \log m + \log \log n) \right\}, \\ &\geq \exp \left\{ -\frac{\varepsilon^2 (1 + \delta)^2 (1 + \gamma)}{\sigma^2 (1 - \delta)} \cdot (\log \log m + \log \log n) \right\}, \end{aligned} \right. \end{aligned}$$

for n large and any $\gamma > 0$.

For the upper bound we define, with some constant $c > 0$ to be defined later,

$$(m_i, n_j) = (e^{c i / \log(i+1)}, e^{c j / \log(j+1)}) \quad \text{for } i, j \geq \max\{\log(1/c)/c, 1\},$$

to obtain

$$\begin{aligned} P\left(T'_{(m_i, n_j)}, (m_i + m_i / \log \log m_i, n_j + n_j / \log \log n_j) > \varepsilon \sqrt{2m_i n_j \frac{\log \log m_i + \log \log n_j}{\log \log m_i \log \log n_j}}\right) \\ \leq C \left(\frac{\log i \log j}{ij} \right)^{-\frac{\varepsilon^2 (1 - \delta)^3}{\sigma^2}}. \end{aligned}$$

The analog of (4.6) turns into

$$P''(m, n) \leq C \left(\frac{\log \log m + \log \log n}{\log(mn)} \right)^N,$$

so that, along the current subset, we obtain

$$P''(m_i, n_j) \leq C \left(\frac{(\log i + \log j) \log i \log j}{i + j} \right)^N,$$

and therefore, in complete analogy with the previous case (use $N \geq 3$ here),

$$\limsup_{i,j \rightarrow \infty} \frac{|T''_{(m_i, n_j), (m_i + m_i / \log \log m_i, n_j + n_j / \log \log n_j)}|}{\sqrt{f_{m_i, n_j}}} \leq \delta \quad \text{a.s.}$$

The argument for T''' is the same as before, so that, recall (3.6),

$$\sum_{m,n} P(|X_{m,n}| > \eta \sqrt{f_{m,n}}) = \sum_{m,n} P(|X| > \eta \sqrt{2mn \frac{\log \log m + \log \log n}{\log \log m \log \log n}}) < \infty,$$

since, by assumption, $E X^2 \log^+ |X| \log^+ \log^+ |X| < \infty$.

Finally, by combining the pieces, it follows that

$$\limsup_{i,j \rightarrow \infty} \frac{T_{(m_i, n_j), (m_i + m_i / \log \log m_i, n_j + n_j / \log \log n_j)}}{\sqrt{2f_{m_i, n_j}}} \leq \sigma(1 - \delta)^{-3/2} + \delta \quad \text{a.s.}, \quad (4.13)$$

for $(m_i, n_j) = (e^{ci/\log(i)}, e^{cj/\log(j)})$ with any $c > 0$.

Sufficiency for the entire field

Once again we have proved the theorem for a suitable subset, and it remains to show that our process behaves accordingly for the entire field. In order to achieve this we use the same procedure as for the corresponding part of the proof of part (i) with logarithms replaced by iterated logarithms.

We first note that here,

$$\begin{aligned} \frac{m_{i+1} - m_i}{a_{m_i}} &\leq \frac{\exp\left(\frac{c(i+1)}{\log(i+1)}\right) - \exp\left(\frac{ci}{\log(i+1)}\right)}{\exp\left(\frac{ci}{\log(i+1)}\right)} \log(ci/\log(i+1)) \leq 2c, \\ \frac{a_{m_{i+1}}}{a_{m_i}} &\leq \left(\frac{\exp\left(\frac{c(i+1)}{\log(i+1)}\right)}{\exp\left(\frac{ci}{\log(i+1)}\right)} \right) \frac{\log(ci/\log(i+1))}{\log(c(i+1)/\log(i+2))} \leq 1 + 2c, \end{aligned}$$

for all integers i considered, so that, by choosing $c = \eta^2/2$, the analog of (4.12) becomes

$$\begin{cases} \text{Var}(T_{(m_i, n_j), (m_{i+1}, n_{j+1})}) \leq \sigma^2 \eta^4 a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1} + a_{m_{i+1}}^{(1)}, n_{j+1})}) \leq \sigma^2 (1 + 2\eta^2) \eta^2 a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1}, n_{j+1} + a_{n_{j+1}}^{(2)})}) \leq \sigma^2 (1 + 2\eta^2) \eta^2 a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1} + a_{m_{i+1}}^{(1)}, n_{j+1} + a_{n_{j+1}}^{(2)})}) \leq \sigma^2 (1 + 2\eta^2)^2 a_{m_i, n_j}, \end{cases} \quad (4.14)$$

for $i, j \geq i_0$.

We omit the remaining details.

Lower bound

Let $c > 1$ and define $m_i = n_i = \exp\{ci/\log(i+1)\}$. Now,

$$\begin{aligned} \frac{e^{ci/\log(i+1)} + e^{ci/\log(i+1)}/\log(ci/\log(i+1))}{e^{c(i+1)/\log(i+2)}} &= \frac{1 + 1/\log(ci/\log(i+1))}{\exp(c(i+1)/\log\{(i+1)(1 + 1/(i+1))\})} \\ &\sim \frac{1 + 1/\log(ci)}{\exp(c/\log(i+1))} < 1, \end{aligned}$$

eventually, which, in analogy to above tells us that the windows along the subset (m_i, n_j) , for i, j large, are disjoint, that is, that different blocks consist of independent random variables. The lower exponential bound in (4.1) then takes care of T' , after which the conclusion of this part of the proof is analogous to the conclusion of the lower bound for part (i) of the theorem.

Necessity

In this case

$$\limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{f_{m,n}}} = \limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{mn \frac{\log \log m + \log \log n}{\log \log m \log \log n}}} < \infty \quad \text{a.s.},$$

which, in view of (3.7), is equivalent $E X^2 \log^+ |X| \log^+ \log^+ |X| < \infty$. The remaining part follows as before.

4.3 Proof of Theorem 2.1(iii)**Sufficiency**

The proof follows the same pattern as that of the previous ones, although the rates of the stretches of the windows now differ in the two directions. Accordingly, the components of the subset will grow at different rates. We have

$$a_{m,n} = \frac{mn}{\log m \log \log n}, \quad d_{m,n} = 2(\log \log m + \log \log n), \quad f_{m,n} = \frac{2mn (\log \log m + \log \log n)}{\log m \log \log n}.$$

The Kolmogorov exponential upper bound along the subset $(m_i, n_j) = (e^{\sqrt{ci}}, e^{\sqrt{cj}})$, $i, j \geq \max\{1/c, 1\}$ with an arbitrarily small $c > 0$, then tells us that

$$P\left(T'_{(m_i, n_j)}, (m_i + m_i / \log m_i, n_j + n_j / \log \log n_j) > \varepsilon \sqrt{2f_{m_i, n_j}}\right) \leq C(ij)^{-\frac{\varepsilon^2(1-\delta)^3}{\sigma^2}}.$$

As for P'' we obtain

$$P''(m_i, n_j) \leq C \left(\frac{\log \sqrt{i} + \log \sqrt{j}}{(\sqrt{i} + \sqrt{j})^2} \right)^N,$$

and, hence, that for $N \geq 3$,

$$\limsup_{i,j \rightarrow \infty} \frac{|T''_{(m_i, n_j)}, (m_i + m_i / \log m_i, n_j + n_j / \log \log n_j)|}{\sqrt{f_{m_i, n_j}}} \leq \delta \quad \text{a.s.},$$

and for T''' , via (3.8) and the fact that $E X^2 (\log^+ |X|)^2 < \infty$,

$$\sum_{m,n} P(|X_{m,n}| > \eta \sqrt{f_{m,n}}) = \sum_{m,n} P\left(|X| > \eta \sqrt{\frac{mn(\log \log m + \log \log n)}{\log m \log \log n}}\right) < \infty.$$

Combining everything finally yields

$$\limsup_{i,j \rightarrow \infty} \frac{T_{(m_i, n_j)}, (m_i + m_i / \log m_i, n_j + n_j / \log \log n_j)}{\sqrt{2f_{m_i, n_j}}} \leq \sigma(1 - \delta)^{-3/2} + \delta \quad \text{a.s.} \quad (4.15)$$

Sufficiency for the entire field

This case is similar to the first one. Since the subsequence for the n -coordinate is denser than in the $\log \log$ -case in part (ii), the task to fill the gaps is even simpler than before.

Lower bound

Once again we let $c > 2$ and set $m_i = n_i = e^{\sqrt{ci}}$. For the first coordinate we have $m_i + m_i / \log m_i < m_{i+1}$ eventually, i.e., the blocks consist of independent random variables. The lower exponential bound now yields

$$\sum_{i,j} P\left(T'_{(m_i, n_j)}, (m_i + m_i / \log m_i, n_j + n_j / \log \log n_j) > (1 - \eta) \sigma \sqrt{4 \frac{mn(\log \log m_i + \log \log n_j)}{\log m_i \log \log n_j}}\right) = \infty,$$

and the lower bound follows as before.

Necessity

This time

$$\limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{f_{m,n}}} = \limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{\frac{2mn(\log \log m + \log \log n)}{\log m \log \log n}}} < \infty \quad \text{a.s.,}$$

and the necessity follows along the usual lines.

5 Additional results

In this section we consider windows of the form

$$T_{(m,n), (m+m^\alpha, n+n/\log n)},$$

that is, the windows are rectangles where one side has a length $a_n^{(2)} = n/\log n$ as before, whereas the other one has length $a_n^{(1)} = m^\alpha$ for some $\alpha \in (0, 1)$, and, hence, is *much* shorter, so that, noticing that $r(n) = (1 - \alpha) \log n$, our usual quantities become

$$a_{m,n} = \frac{m^\alpha n}{\log n}, \quad d_{m,n} = (1 - \alpha)(\log m + \log n), \quad f_{m,n} = (1 - \alpha) \frac{m^\alpha n (\log m + \log n)}{\log n}.$$

Here is now our result for this setting.

Theorem 5.1 *Let $0 < \alpha < 1$. If*

$$E X^{2/\alpha} (\log^+ |X|)^{-1/\alpha} < \infty, \quad \text{and} \quad E(X) = 0, \quad E(X^2) = \sigma^2,$$

then

$$\limsup_{m,n \rightarrow \infty} \frac{T_{(m,n), (m+m^\alpha, n+n/\log n)}}{\sqrt{2m^\alpha n \frac{(1-\alpha) \log(mn)}{\log n}}} = \sigma \quad \text{a.s.}$$

Conversely, if

$$P\left(\limsup_{m,n \rightarrow \infty} \frac{|T_{(m,n), (m+m^\alpha, n+n/\log n)}|}{\sqrt{m^\alpha n \frac{\log(mn)}{\log n}}} < \infty\right) > 0,$$

then $E X^{2/\alpha} (\log^+ |X|)^{-1/\alpha} < \infty$ and $E(X) = 0$.

PROOF. As for the moment condition,

$$\begin{aligned} \limsup_{m,n \rightarrow \infty} \frac{|X_{m,n}|}{\sqrt{m^\alpha n \frac{\log(mn)}{\log n}}} < \infty \quad \text{a.s.} &\iff \sum_{m,n} P\left(|X| > \sqrt{m^\alpha n \frac{\log(mn)}{\log n}}\right) < \infty \\ &\iff E(X^2 / \log^+ |X|)^{1/\alpha} < \infty, \end{aligned} \quad (5.1)$$

which will be verified in the Appendix.

The next step is to truncate and split the window into the usual three parts. Toward that end we note that (3.1) becomes

$$b_{m,n} = \frac{\sigma \delta}{\varepsilon} \sqrt{\frac{m^\alpha n}{(1 - \alpha)(\log m \log n + (\log n)^2)}}, \quad (5.2)$$

after which we introduce the standard three components $X'_{m,n}$, $X''_{m,n}$, and $X'''_{m,n}$ of the summand $X_{m,n}$ and observe that

$$\frac{a_{m,n}}{b_{m,n}} = \frac{\sigma \delta}{\varepsilon} \sqrt{f_{m,n}},$$

so that, via the usual estimates,

$$|E T'_{(m,n), (m+m^\alpha, n+n/\log n)}| = o(\sqrt{f_{m,n}}) \quad \text{as } m, n \rightarrow \infty.$$

With this in mind the exponential bounds yield

$$P\left(T'_{(m,n), (m+m^\alpha, n+n/\log n)} > \varepsilon \sqrt{2f_{m,n}}\right) \begin{cases} \leq \exp\left\{-\frac{\varepsilon^2(1-\delta)^3}{\sigma^2} \cdot (1-\alpha) \log(mn)\right\}, \\ \geq \exp\left\{-\frac{\varepsilon^2(1+\delta)^2(1+\gamma)}{\sigma^2(1-\delta)} \cdot (1-\alpha) \log(mn)\right\}, \end{cases} \quad (5.3)$$

for any $\gamma > 0$ and $\max\{m, n\}$ large.

We note, in passing, that the sum of the probabilities converges for suitably chosen geometrically increasing sequences.

As for T'' , (4.5) turns into

$$\begin{aligned} P(|T''_{(m,n), (m+m^\alpha, n+n/\log n)}| > \eta \sqrt{f_{m,n}}) &\leq C \left(\frac{m^\alpha n}{\log n}\right)^N \left(\frac{E|X|^{2/\alpha}}{\left(\frac{m^\alpha n}{(\log n)^2 + \log m \log n}\right)^{1/\alpha}}\right)^N \\ &= C \frac{((\log n)^{1-\alpha} (\log n + \log m))^{N/\alpha}}{m^{(1-\alpha)N} n^{((1/\alpha)-1)N}}, \end{aligned}$$

so that

$$\sum_{m,n} P(|T''_{(m,n), (m+m^\alpha, n+n/\log n)}| > \eta \sqrt{f_{m,n}}) < \infty \quad (5.4)$$

whenever $N > \max\{1/(1-\alpha), \alpha/(1-\alpha)\} = 1/(1-\alpha)$, that is (cf. [8]), recalling that $N\delta > \eta$ as before,

$$\limsup_{m,n \rightarrow \infty} \frac{|T''_{(m,n), (m+m^\alpha, n+n/\log n)}|}{\sqrt{f_{m,n}}} \leq \frac{\delta}{1-\alpha}.$$

By copying the arguments for T''' from above (as well as from our predecessors) it finally follows that

$$\frac{T'''_{(m,n), (m+m^\alpha, n+n/\log n)}}{\sqrt{f_{m,n}}} \xrightarrow{a.s.} 0 \quad \text{as } m, n \rightarrow \infty.$$

Upper bound

For the upper bound, let $c > 0$ and consider the subset

$$(m_i, n_j) = \left(c i^{1/(1-\alpha)}, c j^{1/(1-\alpha)}\right) \quad \text{for } i, j \geq \max\{c^{\alpha-1}, 1\}. \quad (5.5)$$

Summarizing the three contributions, and taking the arbitrariness of δ and η into account, we have shown that for this subset we have

$$\limsup_{i,j \rightarrow \infty} \frac{T_{(m_i, n_j), (m_i+m_i^\alpha, n_j+n_j/\log n_j)}}{\sqrt{2f_{m_i, n_j}}} \leq \sigma \quad \text{a.s.} \quad (5.6)$$

for any $c > 0$.

Sufficiency for the entire field

Here we have that for $i, j \geq c^{2(\alpha-1)}$ and $\log j/j \leq 1/\sqrt{j}$ with $c > 0$, sufficiently small,

$$\left\{ \begin{aligned} m_{i+1} - m_i &\sim \frac{c}{1-\alpha} i^{\alpha/(1-\alpha)} = \frac{c^{1-\alpha}}{1-\alpha} a_{m_i}^{(1)}, \\ n_{j+1} - n_j &\sim \frac{c}{1-\alpha} j^{\alpha/(1-\alpha)} = \frac{\log j}{(1-\alpha)^2 j} a_{n_j}^{(2)} \leq \frac{c^{1-\alpha}}{(1-\alpha)^2} a_{n_j}^{(2)}, \\ a_{m_{i+1}}^{(1)} / a_{m_i}^{(1)} &\sim 1 + \frac{\alpha}{1-\alpha} \cdot \frac{1}{i} \leq 1 + \frac{\alpha c^{2(1-\alpha)}}{1-\alpha}, \\ a_{n_{j+1}}^{(2)} / a_{n_j}^{(2)} &= \left(1 + \frac{1}{j}\right)^{1/(1-\alpha)} \cdot \frac{\log j}{\log(j+1)} \leq 1 + \frac{1}{1-\alpha} \cdot \frac{1}{j} \leq 1 + \frac{c^{2(1-\alpha)}}{1-\alpha}. \end{aligned} \right. \quad (5.7)$$

By choosing $\eta^2 = c^{1-\alpha}/(1-\alpha)^2$ (which implies that $c = ((1-\alpha)^2\eta^2)^{1/(1-\alpha)} \leq \eta^2$), the estimates for the variances turn out as follows:

$$\left\{ \begin{array}{l} \text{Var}(T_{(m_i, n_j), (m_{i+1}, n_{j+1})}) \leq \sigma^2 \frac{c^{2(1-\alpha)}}{(1-\alpha)^3} a_{m_i, n_j} \leq \sigma^2 \eta^4 a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1} + a_{m_{i+1}}^{(1)}, n_{j+1})}) \leq \sigma^2 \left(1 + \frac{c^{1-\alpha} + \alpha c^{2(1-\alpha)}}{1-\alpha}\right) \frac{c^{1-\alpha}}{(1-\alpha)^2} a_{m_i, n_j} \\ \leq \sigma^2 \left(1 + \frac{2c^{1-\alpha}}{(1-\alpha)^2}\right) \frac{c^{1-\alpha}}{(1-\alpha)^2} a_{m_i, n_j} \\ \leq \sigma^2 (1 + \eta^2) \eta^2 a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1}, n_{j+1} + a_{n_{j+1}}^{(2)})}) \leq \sigma^2 \frac{c^{1-\alpha}}{1-\alpha} \left(\frac{c^{1-\alpha}}{(1-\alpha)^2} + 1 + \frac{c^{2(1-\alpha)}}{1-\alpha}\right) a_{m_i, n_j} \\ \leq \sigma^2 \frac{c^{1-\alpha}}{1-\alpha} \left(1 + \frac{2c^{1-\alpha}}{(1-\alpha)^2}\right) a_{m_i, n_j} \\ \leq \sigma^2 \eta^2 (1 + 2\eta^2) a_{m_i, n_j}, \\ \text{Var}(T_{(m_i, n_j), (m_{i+1} + a_{m_{i+1}}^{(1)}, n_{j+1} + a_{n_{j+1}}^{(2)})}) \leq \sigma^2 \left(1 + \frac{c^{1-\alpha} + \alpha c^{2(1-\alpha)}}{1-\alpha}\right) \\ \times \left(\frac{c^{1-\alpha}}{(1-\alpha)^2} + 1 + \frac{c^{2(1-\alpha)}}{1-\alpha}\right) a_{m_i, n_j} \\ \leq \sigma^2 \left(1 + \frac{2c^{1-\alpha}}{(1-\alpha)^2}\right)^2 a_{m_i, n_j} \\ \leq \sigma^2 (1 + 2\eta^2)^2 a_{m_i, n_j}, \end{array} \right. \quad (5.8)$$

for $i, j \geq i_0$.

The remaining part of the proof follows the usual procedure with obvious changes and is omitted.

Lower bound

For the lower bound we set, with some $c > 1$,

$$m_i = n_i = c i^{1/(1-\alpha)},$$

so that combining the contributions above, taking the arbitrariness of δ , γ , and η into account, we obtain

$$\limsup_{i, j \rightarrow \infty} \frac{T_{(m_i, n_j), (m_i + m_i^\alpha, n_j + n_j / \log n_j)}}{\sqrt{2f_{m_i, n_j}}} \geq \sigma \quad \text{a.s.} \quad (5.9)$$

Namely, since, for the current subset we have $m_i + m_i^\alpha < m_{i+1}$ eventually, the windows of the subset are disjoint, so that (5.9) follows in view of the second Borel–Cantelli lemma, after which, once again, the fact that the limsup over *all* windows is at least as large as the limsup over a subset establishes (5.9).

Necessity

This part follows the standard pattern.

6 Some final remarks

1. The subsequences we have used in the proof of the upper bound have, throughout been the same for both coordinates, even in those cases when the edges of the windows grow at different rates, viz., in the proofs of Theorem 2.1(iii) and (5.5). Intuitively one might imagine subsequences that grow at different rates to be more natural. However, since there is less cancellation going on along the direction of the shorter edge, that is, since the fluctuations along the direction of the shorter edge are stronger, it turns out that the shortest edge determines the moment condition

(as in [9]). Our choice of subset implies that the windows of the subset are “covered” many times along the longer direction, which, on the one hand is not necessary, but on the other does not exhibit any “harmful waste”, in that our estimates produce the best result.

2. Analogous to our previous results [8, 9] we can obtain limit theorems over subsets, which, in particular, show that all reals between the limit inferior and the limit superior are limit points.
3. An analysis of the proofs shows that it is also possible to formulate a more general result. However, the proof is more tedious and the moment condition is not so explicit.

Assume that $L_i(x) \nearrow \infty$ for $i = 1, 2$ are slowly varying functions satisfying (1.2), and suppose that $L_1(x) \geq L_2(x)$ for $x > \text{some } x_0$. Further, define $r(n) = \log(L_1(n) \log n)$ and

$$G(u, v) = \frac{uv}{L_1(u) L_2(v)} (r(u) + r(v))$$

with the associated M -function

$$M(x) = |\{(u, v) \in [1, \infty) \times [1, \infty) : G(u, v) \leq x\}|.$$

Then the following result holds:

Theorem 6.1 *If $EM(X^2) < \infty$, $E(X) = 0$, and $E(X^2) = \sigma^2$, then*

$$\limsup_{m, n \rightarrow \infty} \frac{T_{(m, n), (m+m/L_1(m), n+n/L_2(n))}}{\sqrt{2G(m, n)}} = \sigma \quad a.s. \quad (6.1)$$

Conversely, if the lim sup is finite with positive probability, then $EM(X^2) < \infty$, $E(X) = 0$, and (6.1) holds with $\text{Var } X = \sigma^2$.

REMARK 6.1 Note that $M(x)/x \rightarrow \infty$, that is, the second moment exists if $EM(X^2) < \infty$.

REMARK 6.2 Due to the generality it is not possible to describe the function $M(x)$ more precisely, but one can prove (see the Appendix) that

$$M(x) \leq C \frac{x L_2(x)}{r(x)} \int_0^{L_1(x)/r(x)} \frac{L_1(u)}{u} du. \quad (6.2)$$

The proof follows the lines of the proofs above using the techniques from [7] with subsequences $m_k = n_k = \psi(ck)$ where $\psi(x) = \varphi^{-1}(x)$ with $\varphi(x) = \int^x L_1(u)/u du \nearrow \infty$ and being in $\mathcal{SV}$.

REMARK 6.3 For Theorem 2.1 the estimate in (6.2) is precise enough to give the correct moment assumptions. $\square$

Appendix

Proofs of (3.6) – (3.8), (5.1) and (6.2)

In this appendix we present the “elementary but tedious calculations” needed for the verification of the second half of Corollary 3.1 and the further analogs. Throughout it is tacitly assumed that all logarithms and iterated logarithms are to be interpreted as $\max\{\log x, 1\}$ and $\max\{\log \log x, 1\}$, respectively. Moreover, we shall use $a(x) \approx b(x)$ to denote that $0 < c_1 \leq a(x)/b(x) \leq c_2$. Finally, the constant C below may be different at each occurrence.

Proof of (3.6)

Since $\log u/u \searrow$ we obtain

$$\begin{aligned} I_1(u) &= \iint_{xy(\log \log x + \log \log y)/\log x \log y \leq u} 1 \, dx dy = \iint_{xy(\log(\log x \log y))/\log x \log y \leq u} 1 \, dx dy \\ &\leq \iint_{xy \leq u(\log u)^2/\log((\log u)^2)} 1 \, dx dy \leq C \iint_{xy \leq u(\log u)^2/\log \log u} 1 \, dx dy \leq Cu(\log u)^3/\log \log u, \end{aligned}$$

and

$$\begin{aligned} I_1(u) &\geq \int_{\sqrt{u}}^u \int_{x \geq y^{-1}u(\log \sqrt{u})^2 / \log((\log \sqrt{u})^2)} 1 \, dx dy \geq C \int_{\sqrt{u}}^u \int_{x \leq (4y)^{-1}u(\log u)^2 / \log \log u} 1 \, dx dy \\ &\geq Cu(\log u)^3 / \log \log u. \end{aligned}$$

This shows that $I_1(u) \approx u(\log u)^3 / \log \log u$ as $u \rightarrow \infty$, i.e., we may choose $M(x) = x(\log x)^3 / \log \log x$, and (3.6) follows.

Proof of (3.7)

We have

$$\begin{aligned} I_2(u) &= \iint_{xy \log(\log x \cdot \log y) / \log \log x \log \log y \leq u} 1 \, dx dy = \iint_{xy(\frac{1}{\log \log x} + \frac{1}{\log \log y}) \leq u} 1 \, dx dy \\ &\leq \iint_{xy \leq 2u \log \log u} 1 \, dx dy \leq Cu \log u \log \log u, \end{aligned}$$

and

$$I_2(u) \geq \int_{\sqrt{u}}^u \int_{x \leq 2y^{-1}u \log \log \sqrt{u}} 1 \, dx dy \geq Cu \log u \log \log u.$$

This shows that $I_2(u) \approx \log u \log \log u$ as $u \rightarrow \infty$, from which we conclude that (3.7) holds.

Proof of (3.8)

We first note that, $\frac{\log \log x}{\log \log y} + 1 \geq 1$, and that for $\sqrt{u} \leq x, y \leq u$,

$$\frac{\log \log x}{\log \log y} + 1 \leq \frac{\log \log u}{\log \log \sqrt{u}} + 1 = \frac{\log \log u}{-\log 2 + \log \log u} + 1 = 2 + \frac{\log 2}{-\log 2 + \log \log u} \leq C,$$

so that, on the one hand,

$$\begin{aligned} I_3(u) &= \iint_{xy(\log \log x + \log \log y) / \log x \log \log y \leq u} 1 \, dx dy = \iint_{\frac{xy}{\log x} (\frac{\log \log x}{\log \log y} + 1) \leq u} 1 \, dx dy \\ &\leq \iint_{\frac{xy}{\log x} \leq u} 1 \, dx dy \leq \int_1^{u \log u} u \frac{\log x}{x} dx = \frac{u(\log u + \log \log u)^2}{2}, \end{aligned}$$

and on the other,

$$I_3(u) \geq \iint_{\frac{Cxy}{\log x} \leq u} 1 \, dx dy \sim Cu(\log u)^2.$$

This shows that $I_3(u) \approx u(\log u)^2$ as $u \rightarrow \infty$, which proves (3.8).

Proof of (5.1)

We first note that, since $1 < x, y < u$, we have

$$x^\alpha y \frac{\log x}{\log y} \leq x^\alpha y \left(\frac{\log x}{\log y} + 1 \right) = x^\alpha y \frac{\log x + \log y}{\log y} \leq 2x^\alpha y \frac{\log u}{\log y}.$$

Thus,

$$\begin{aligned}
I_4(u) &= \iint_{x^\alpha y (\frac{\log x}{\log y} + 1) \leq u} 1 \, dx dy \leq \iint_{x^\alpha y \frac{\log x}{\log y} \leq u} 1 \, dx dy = \iint_{\frac{y}{\log y} \leq \frac{u}{x^\alpha \log x}} 1 \, dx dy \\
&\leq C \int_1^{(u/\log u)^{1/\alpha}} \int_{y \leq \frac{u \log u}{x^\alpha \log x} - \frac{\alpha u}{x^\alpha} - \frac{u \log \log x}{x^\alpha \log x}} 1 \, dy dx \\
&= C \int_1^{(u/\log u)^{1/\alpha}} \left(\frac{u \log u}{x^\alpha \log x} - \frac{\alpha u}{x^\alpha} - \frac{u \log \log x}{x^\alpha \log x} \right) dx \\
&= C \left(\frac{u \log u}{1-\alpha} \left\{ \left[\frac{x^{1-\alpha}}{\log x} \right]^{(u/\log u)^{1/\alpha}} + \int_1^{(u/\log u)^{1/\alpha}} \frac{1}{x^\alpha (\log x)^2} dx \right\} \right. \\
&\quad \left. - \frac{\alpha u}{1-\alpha} \left(\frac{u}{\log u} \right)^{(1/\alpha)(1-\alpha)} - \frac{u}{1-\alpha} \left(\frac{u}{\log u} \right)^{1-1/\alpha} \frac{\alpha \log \log u}{\log u} \left(1 + \mathcal{O} \left(\frac{\log \log u}{\log u} \right) \right) \right) \\
&\leq C \left(\frac{u \log u}{1-\alpha} \left\{ \frac{(u/\log u)^{(1/\alpha)(1-\alpha)}}{\frac{1}{\alpha} (\log u - \log \log u)} + \tilde{C} \left[\frac{1}{1-\alpha} \cdot \frac{x^{1-\alpha}}{(\log x)^2} \right]^{(u/\log u)^{1/\alpha}} \right\} \right. \\
&\quad \left. - \frac{\alpha u}{1-\alpha} \left(\frac{u}{\log u} \right)^{(1/\alpha)(1-\alpha)} - \frac{u}{1-\alpha} \left(\frac{u}{\log u} \right)^{1-1/\alpha} \frac{\alpha \log \log u}{\log u} \right) \\
&\leq C u \log u \frac{(u/\log u)^{(1/\alpha)(1-\alpha)}}{(\log u)^2} = C \left(\frac{u}{\log u} \right)^{1/\alpha}.
\end{aligned}$$

As for the lower bound,

$$\begin{aligned}
I_4(u) &= \iint_{x^\alpha y \frac{\log x + \log y}{\log y} \leq u} 1 \, dx dy \geq \iint_{x^\alpha y \frac{\log u + \log u}{\log y} \leq u} 1 \, dx dy = \int^{u/\log u} \int_{x \leq \left(\frac{u \log y}{2y \log u} \right)^{1/\alpha}} 1 \, dx dy \\
&= \left(\frac{u}{\log u} \right)^{1/\alpha} \int^{u/\log u} \left(\frac{\log y}{2y} \right)^{1/\alpha} dy \geq C \left(\frac{u}{\log u} \right)^{1/\alpha},
\end{aligned}$$

since the last integral has a limit as $u \rightarrow \infty$. Hence, $I_4(u) \approx (u/\log u)^{1/\alpha}$ and $M(x) = (x/\log x)^{1/\alpha}$.

Proof of (6.2)

For the general case the corresponding integral can be estimated from above by

$$\begin{aligned}
I_5(u) &= \int \int_{\frac{xy(r(x)+r(y))}{L_1(x)L_2(y)} \leq u} 1 \, dx dy \leq \int \int_{\frac{y}{L_2(y)} \leq u \frac{L_1(x)}{x r(x)}} 1 \, dy dx \\
&\leq C u L_2(u) \int^{u L_1(u)/r(u)} \frac{L_1(x)}{x r(x)} dx \leq C u \frac{L_2(u)}{r(u)} \int^{u L_1(u)/r(u)} \frac{L_1(x)}{x} dx \\
&\leq C u \frac{L_2(u)}{r(u)} \int^{u L_1(u)/r(u)} \frac{L_1(x)}{x} dx,
\end{aligned}$$

where we used [7], Lemma 3.1, for the second to last inequality.

Interchanging the roles of the two variables will in general lead to a poorer estimate.

In the cases discussed in detail the estimates are precise enough, since the lower bound

$$\begin{aligned}
I_5(u) &= \int \int_{\frac{xy(r(x)+r(y))}{L_1(x)L_2(y)} \leq u} 1 \, dx dy \geq \int_{u^\beta}^{c u L_1(u)/r(u)} \int_{\frac{xy(r(u)+r(y))}{L_1(u^\beta)L_2(y)} \leq u} 1 \, dx dy \\
&= \int_{u^\beta}^{c u L_1(u)/r(u)} \int_{y \leq \frac{u L_2(u^\beta)}{2r(u)} \frac{L_1(x)}{x}} 1 \, dy dx \geq C \frac{u L_2(u^\beta)}{r(u)} \int_{u^\beta}^{u L_1(u)/r(u)} \frac{L_1(x)}{x} dx
\end{aligned}$$

e.g. for $\beta = 1/2$, leads to the same functions (up to constants), that is, we rediscover our results in the examples discussed above.

Finally, since the last integral on the right-hand side increases faster than the logarithm and since $r(u) = o(\log u)$, it follows that $M(u)/u \rightarrow \infty$.

References

- [1] BINGHAM, N. H., GOLDIE, C. M. AND TEUGELS, J.L. (1987). *Regular Variation*. Cambridge University Press, Cambridge.
- [2] CHOW, Y.S. AND LAI, T.L. (1975). Some one-sided theorems on the tail distribution of sample sums with applications to the last exit time and largest excess of boundary crossings. *Trans. Amer. Math. Soc.* **208**, 51-72.
- [3] CSÖRGŐ, M. AND RÉVÉSZ, P. (1981). *Strong Approximations in Probability and Statistics*, Academic Press, New York.
- [4] ERDŐS, P. AND RÉNYI, S. (1970). On a new law of large numbers. *J. Analyse Math.* **23**, 103-111.
- [5] FELLER, W. (1968). An extension of the law of the iterated logarithm to variables without variance. *J. Math. Mech.* **18**, 343-355.
- [6] GUT, A. (2007). *Probability: A Graduate Course*, Corr. 2nd printing. Springer-Verlag, New York.
- [7] GUT, A., JONSSON, F. AND STADTMÜLLER, U. (2009). Between the LIL and LSL. *Bernoulli* (to appear).
- [8] GUT, A. AND STADTMÜLLER, U. (2008a). Laws of the single logarithm for delayed sums of random fields. *Bernoulli*, **14**, 249-276.
- [9] GUT, A. AND STADTMÜLLER, U. (2008b). Laws of the single logarithm for delayed sums of random fields II. *J. Math. Anal. Appl.* **346**, 403-413.
- [10] HARTMAN, P. AND WINTNER, A. (1941). On the law of the iterated logarithm. *Amer. J. Math.* **63**, 169-176.
- [11] LAI, T.L. (1974). Limit theorems for delayed sums. *Ann. Probab.* **2**, 432-440.
- [12] MARTIKAINEN, A.I. (1980). A converse to the law of the iterated logarithm for a random walk. *Theory Probab. Appl.* **25**, 361-362.
- [13] PARANJAPE, S.R. AND PARK C. (1973). Laws of iterated logarithm of multiparameter Wiener processes. *J. Multivariate Anal.* **3**, 132-136.
- [14] PRUITT, W.E. (1981). General one-sided laws of the iterated logarithm. *Ann. Probab.* **9**, 1-48.
- [15] ROSALSKY, A. (1980). On the converse to the iterated logarithm law. *Sankhyā, Ser. A* **42**, 103-108.
- [16] STRASSEN, V. (1966). A converse to the law of the iterated logarithm. *Z. Wahrsch. verw. Gebiete* **4**, 265-268.
- [17] WICHURA, M.J. (1973) Some Strassen-type laws of the iterated logarithm for multiparameter stochastic processes with independent increments. *Ann. Probab.* **1**, 272-296.

Allan Gut, Department of Mathematics, Uppsala University, Box 480, SE-751 06 Uppsala, Sweden;

Email: allan.gut@math.uu.se

URL: <http://www.math.uu.se/~allan>

Ulrich Stadtmüller, Ulm University, Department of Number Theory and Probability Theory,
D-89069 Ulm, Germany;

Email ulrich.stadtmueller@uni-ulm.de

URL: <http://www.mathematik.uni-ulm.de/matheIII/members/stadtmueller/stadtmueller.html>